

Sketch map showing location of the Middle and Upper Devonian sections exposed along Zolotukha and Gryaznukha brooks, SW Rudny Altai, Altai-Sayan Folded Area (ASFA), southwestern Siberia.

associations come from fine-grained pale-gray, greenish-gray, and black siliceous mudstones, shales and cherts from four stratigraphic intervals: uppermost Givetian–lower Frasnian, middle Frasnian, and upper Frasnian.

The first association is found in section SB-031, in the Zolotukha Formation, exposed on the right bank of Zolotukha Brook (Fig. 3, beds 8, 10, 13, 15, 17, 18, 45). Conodonts *Klapperina disparilis* (Ziegler & Klapper) and *Polygnathus ovinodosus* Ziegler & Klapper obtained from bed 9 (Bakharev *et al.* 2004) indicate an uppermost Givetian–lower Frasnian (*disparilis-falsiovalis* zones) age. The radiolarian fauna is dominated by spherical entactiniids and includes *Trilonche davidi* (Hinde), *Tr. minax* (Hinde), *Tr. palimbola* (Foreman), *Tr. cf. inusitata* (Foreman), *Astroentactinia stellata* Nazarov, *A. paronae*

Global Stratigraphic Scale				Conodont Zone	North-East Kuznetsk Basin	Rudny Altai	
System	Series	Stage	Substage		Horizon, Beds	Formation	
Devonian	Upper	Famennian	J	<i>rhomboidea</i>	Pestichetka	Mitikha Beds	Gryaznukha
				<i>crepida</i>		Kosoy Utyos Beds	
				<i>triangularis</i>		Solomino	
		Frasnian	U	<i>linguiformis</i>	Vassiro	Pozharishchevo	Razdol'noe
				<i>rhenana</i>		Strel'naya	Gerikhovskoe
	M		<i>jameiae</i>	Zolotukha			
			<i>hassi</i>				
	Middle	Givetian	U	<i>punctata</i>	Mazalovskiy Kitat	Davydovka	
				<i>transitans</i>			
				<i>falsiovalis</i>			
<i>disparilis</i>							
<i>hermanni-cristatus</i>							

Regional stratigraphic charts for the Middle-Upper Devonian of the Rudny Altai (after Yolkin *et al.* 2005, Weddige 2006).

(Hinde), and *Palaeoscenidium cladophorum* Deflandre (Fig. 4A, E, F).

A second association was revealed from section S-033 (Fig. 3, beds 6, 8, 12, 14, 18) of the Zolotukha Formation, provisionally dated as middle Frasnian (?*hassi* conodont zone), because overlying strata of section S-034 (see Fig. 1) yielded conodonts *Polygnathus uchtensis* Ovnatanova & Kuzmin, *Palmatolepis hassi* Müller & Müller and *Pal. rhenana* Müller. The radiolarian fauna is characterized by spherical *Trilonche davidi* (Hinde), *Tr. minax* (Hinde), *Tr. hindea* (Hinde), *Tr. echinata* (Hinde), *Tr. vetusta* Hinde, *Astroentactinia stellata* Nazarov, and *Palaeoscenidium cladophorum* Deflandre (Fig. 4G).

A third association is from section S-0210 (Fig. 3) in the Gryaznukha Formation. Conodonts *Palmatolepis rhenana* Bischoff, *Pal. jameiae* Ziegler & Sandberg, *Pal. gigas* Müller & Youngquist and ammonoids in this part of the section are dated as upper Frasnian (*rhenana-linguiformis* zones and *Manticoceras Stufen*). Radiolarians are represented by a few specimens of *Trilonche hindea* (Hinde), *Astroentactinia stellata* Nazarov, *Palaeoscenidium cladophorum* Deflandre, and *Palaeoscenidium* sp.


The fourth association is the most diverse. It was recovered from section S-0011 (Fig. 5, beds 10–27) of the upper Frasnian Gryaznukha Formation. It is characterized by *Trilonche guangxiensis* (Li & Wang), *Tr. vetusta* Hinde, *Tr. echinata* (Hinde), *Tr. davidi* (Hinde), *Tr. minax* (Hinde), *Tr. tanheensis* Luo, Aitchison & Wang, *Astroentactinia stellata* Nazarov, *A. vishnevskayae* Afanasieva, *Stigmosphaerostylus* sp. A; *Borisella* cf. *maksimovae* Afanasieva, *Moscovistella* cf. *allbororum* Afanasieva, *Haplentactinia rhinophyusa* Foreman, *Palaeoscenidium cladophorum* Deflandre, *Pal. delicatum* Aitchison, *Pal. tabernaculum* Aitchison, *Ceratoikiscum avimexpectans* Deflandre, *Cer. mirum* Cheng, *Cer. labyrinthum* Cheng,

Cer. delicatum Cheng, *Cer. spinosum* Cheng, *Nazarovites bioculus* Afanasieva, *N. pinnula* Afanasieva, *Cancellientactinia acifera* gen. et sp. nov., and *Palaethalomnus* sp. (Figs 4B–D, H–T, 6A–T). The age of the strata as Upper *rhenana*-*linguiformis* zones has been established by recent findings of conodonts *Palmatolepis* gr. *Pa. delicatula* Branson & Mehl (beds 10, 25), *Pa. cf. Pa. rotunda* Ziegler & Sandberg (beds 18, 27), *Palmatolepis* ssp. (beds 10, 18, 25, 27) and *Polygnathus* sp. (beds 10, 18, 25) (Obut *et al.* 2007). It should be mentioned that in previous reports (Obut 2006), the stratigraphic position of this radiolarian association, was regarded as transitional uppermost Frasnian–Lower Famennian, based on *Cer. avimexpectans* Deflandre, *Cer. mirum* Cheng, *Cer. spinosum* Cheng and *Circuliforma robusta* Cheng, which range into the Famennian–Lower Carboniferous (Cheng 1986, Schwartzapfel & Holdsworth 1996). The other radiolarians present occur in Middle–Upper Devonian sequences worldwide. We also emphasize here that the ranges of many genera and species of Devonian radiolarians require more precise specification.

Radiolarians were extracted from siliceous argillites and cherts of the field area by standard methods using 5–15% hydrofluoric acid, with samples repeatedly washed, and specimens picked under a binocular lens. Detailed observation and photography were made using a Leo 1430 VP SEM. More than 600 specimens in total, mostly slightly broken, were studied. The described collection (registered as # 1099) is stored in the Central Siberian Geological Museum (CGSM) in the Institute of Geology and Mineralogy, Siberian Branch of Russian Academy of Sciences, Novosibirsk, Russia. All figured and described radiolarians come from samples collected from sections SB-031, S-033, S-0210 and S-0011, which crop out in the SW Rudny Altai, near Gornyak Settlement, in the watershed of the Gryaznukha and Zolotukha brooks (Fig. 1).

Radiolarians described below include a new genus and species, as well as some abundant taxa that were not found previously in the sequences examined earlier in this study.

Upper Devonian sections exposed along the Zolotukha and Gryaznukha Brooks, near former Razdol'noe Village, SW Rudny Altai and ranges of key radiolarian taxa. Abbreviation: A – Astroentactinia, T – Trilonche.



- rhyolites
- lavaclastic rhyolites
- siltstones
- siliceous mudstones
- cherty rocks
- clasts of mudstone-siliceous rocks
- olistostromal breccia
- tuffs
- sandstones
- not exposed
- limestones
- conodonts, ammonoids

Family Polyentactiniidae Nazarov, 1974

Genus *Cancellientactinia* gen. nov.

Etymology. – From *cancelli* lat. – lattice and *Entactinia*.

Type species. – *Cancellientactinia acifera* sp. nov., Upper Devonian, upper Frasnian Gryaznukha Formation, Rudny Altai, south of West Siberia, Russia.

Diagnosis. – One spherical latticed shell, constructed by a meshwork of short bars. Inner skeleton represented by an 8-rayed spicule, all eight rays originating from a central point. Eight main more robust spines develop from rays of the internal spicule.

Remarks. – The described genus differs from *Polyentactinia* Foreman (Foreman 1963, p. 281) in having a single internal spicule (without central bar) and the presence of only rod-like spines, and from *Magnisphaera* Won (Won 1997, p. 366) by one latticed shell and an internal spicule.

Occurrence. – Late Devonian, late Frasnian; right bank of Gryaznukha Brook, section S-0011, Rudny Altai, SW Siberia, Russia.

Cancellientactinia acifera sp. nov.

Figure 6A–D

1997 *Polyentactinia* sp.; Aitchison & Stratford, figs 2–12.

Etymology. – From *acifera* lat. – process, thorn.

Holotype. – Complete shell with internal spicule specimen CSGM 1099/1 (Fig. 6A).

Paratypes. – Shells with internal spicule, possessing cross section (specimens CSGM 1099/2, 1099/3).

Type horizon and locality. – Upper Devonian, upper Frasnian Gryaznukha Formation: right bank of Gryaznukha

Brook, section S-0011, bed 25, SW Rudny Altai, SW Siberia, Russia.

Material. – About 20 specimens in CSGM: 2 well preserved shells, 5 damaged shells, more than 10 broken specimens.

Diagnosis. – One robust latticed shell constructed from a meshwork of short bars. Broad rod-like bars enmeshed with fine short thorns and spinules. Eight main spines developed from rays of an internal spicule are rod-like and conical. Test a regular hexagon in cross-section.

Description. – One large sturdy latticed shell is formed from a regular meshwork of short robust rod-like bars. Occurring on the broad bars are specific sculptural elements, fine thorn-like projections of various lengths, sometimes as long as the main spines (Fig. 6B, D). The inner spicule is robust with eight rays developed from a central point. The main spines are rod-like, slightly conical, long, and sometimes equal in length to the test radius. They are decorated with short curved spinules that can sometimes intersect with each other or with the thorns developed on the bars.

Dimensions. – Test diameter 120–180 µm; inner spicule ray diameter 7–10 µm; main spine length 70–90 µm, with base diameter 20 µm; diameter of bars 7–10 µm; by-spine length 10–55 µm.

Occurrence. – The same as for genus.

Family Entactiniidae Riedel, 1967,
emend. Afanasieva, 1999

Genus *Stigmosphaerostylus* Rüst, 1892,
emend. Foreman, 1963

1892 *Stigmosphaerostylus* Rüst; Rüst, p. 142.

1963 *Entactinia* Foreman; Foreman, p. 271.


1988 *Entactinia* Foreman. – Nazarov, pp. 58, 59.

1999 *Stigmosphaerostylus* Rüst. – Braun & Budil, p. 584.

2000a *Entactinia* Foreman. – Afanasieva, p. 41.

Radiolarians from the Zolotukha and Gryaznukha brooks sections, Upper Devonian, Frasnian, SW Rudny Altai. Scale 50 µm.

- A, B – *Astroentactinia stellata* Nazarov; A – section SB-031 (loc. 04090701/20), 1099/10; B – section S-0011 (loc. 00072603/8), 1099/11.
- C, D – *Astroentactinia vishnevskayae* Afanasieva, section S-0011 (loc. 00072603/8); C – 1099/12, D – 1099/13. • E–G – *Trilonche davidi* (Hinde); E, F – section SB-031 (loc. 04090701/20), 1099/20, 1099/21; G – section S-033 (loc. 04090501/4), 1099/22. • H – *Moscovistella allbororum* Afanasieva, section S-0011 (loc. S-0011-10), 1099/56. • I, J – *Trilonche guanxiensis* (Li & Wang), section S-0011 (loc. S-0011-25); I – 1099/31; J – 1099/32.
- K – *Trilonche tangheensis* Luo, Aitchison & Wang, section S-0011 (loc. S-0011-27), 1099/34. • L – *Trilonche echinata* Hinde, section S-0011 (loc. S-0011-10), 1099/25. • M – *Trilonche minax* (Hinde), section S-0011 (loc. S-0011-27), 1099/23. N – *Trilonche hindea* (Hinde), section S-0011 (loc. S-0011-25), 1099/27. • O – *Trilonche vetusta* Hinde, section S-0011 (loc. 00072603/8), 1099/29. • P, S, T – *Stigmosphaerostylus* sp. A, section S-0011 (loc. 00072603/8); P – 1099/41; S – 1099/40; T – 1099/42. • Q, R – *Borisella* cf. *maksimovae* Afanasieva, section S-0011 (loc. 00072603/8); Q – 1099/50; R – 1099/51.



Type species. – *Stigmosphaerostylus notabilis* Rüst, 1892; Lower Carboniferous, Harz Mountains, Germany.

Discussion. – Aitchison & Stratford (1997) regarded the widely-used name *Entactinia* introduced by Foreman (1963) as a junior synonym of *Stigmosphaerostylus* Rüst and proposed Foreman's diagnosis as an amendment. According to the original diagnosis of the genus *Stigmosphaerostylus* by Rüst (1892, p. 142), representatives that possessed one latticed shell with radial spines and an internal spicule constructed of radial rays were assigned to this genus. The description of the only recognized species *St. notabilis* defines the inner structure of this taxon: the inner spicule is constructed of six rays combined in one center (Rüst 1892, taf. X, fig. 2). Foreman's (1963, p. 271) original diagnosis for *Entactinia* is: "A single well-developed latticed or spongy shell (sometimes with external spongy material) and an internal six-rayed double spicule more delicate than the main spines." It should be noted that most of described species in recent literature assigned to *Stigmosphaerostylus* Rüst (*St. ornata* (Hinde), *St. pussila* (Hinde), *St. etheridgei* (Hinde), *St. hystricuosa* (Aitchison), *St. variospina* (Won) are characterized by a six-rayed double internal spicule and one latticed shell (Aitchison & Stratford 1997, Aitchison *et al.* 1999, Braun & Budil 1999, Wang *et al.* 2000, Luo *et al.* 2002).

***Stigmosphaerostylus* sp. A**
Figure 4P, S–T

2004 *Stigmosphaerostylus* sp. A; Umeda *et al.*, pl. III, figs 3, 4.

Type horizon and locality. – Upper Devonian Gryaznukha Formation, right bank of Gryaznukha Brook, section S-0011, loc. S-0011/10, 25, 27, SW Rudny Altai, southwestern Siberia, Russia.

Material. – 10 specimens.

Description. – The one spherical porous shell has six approximately equal main spines that are strongly three-bladed, grooved and tapered at the tip. The shell possesses subcircular pores and a wide interporous area, decorated with fine short thorns. The inner spicule is observed in broken specimens.

Dimensions. – Shell diameter 100–130 µm; main spine length 25–40 µm, with base diameter 15–20 µm.

Remarks. – This form differs from *Stigmosphaerostylus additiva* (Foreman), described from Upper Devonian Hu-

ron Member, Ohio Shale of Ohio, USA (Foreman 1963) and the Frasnian of the South Urals, Russia (Nazarov 1975) in having more delicate main spines, which are smaller in the diameter of their base and length. It resembles *Stigmosphaerostylus bogdanovi* (Afanasieva) reported from the Frasnian and Famennian of the Timan-Pechora region, Russian Platform (Afanasieva 2000a). The form we report has in comparison a larger shell diameter and considerably shorter main spines. Our form possibly represents a new species.

Occurrence. – Upper Devonian, upper Frasnian, Gryaznukha Formation: right bank of Gryaznukha Brook, section S-0011, beds 10, 18, 25, 27, Rudny Altai, southwestern Siberia, Russia.

Genus *Borisella* Afanasieva, 2000

Type species. – *Borisella maksimovae* Afanasieva, 2000; Upper Devonian, Frasnian, Domanic Formation, Timan-Pechora Province, northern Russia.

Diagnosis. – See Afanasieva (2000b, p. 31).

***Borisella* cf. *maksimovae* Afanasieva, 2000**
Figure 4Q, R

2004 *Stigmosphaerostylus* sp. B; Umeda *et al.*, pl. III, fig. 5.


Material. – 5 specimens.

Description. – The one spherical thick porous shell has six rod-like, conical main spines with Y-shape cross-section at the base and a small cavity. The shell possesses subcircular to subangular pores divided by a wide interporous area, framed with fine short thorns, sometimes with additional very short thorns. An inner spicule is observed in broken specimens.

Dimensions. – Shell diameter 100–160 µm; main spine length up to 200 µm, with base diameter 8–10 µm.

Remarks. – Afanasieva (2000b) reported the presence of massive by-spines (sometimes absent) and very long, thin main spines in the material from the Upper Devonian, Frasnian, Domanic Formation of Timan-Pechora Province. The specimens in our collection lack by-spines and the main spines are almost always broken.

Occurrence. – Upper Devonian, upper Frasnian Gryaznukha Formation; right bank of Gryaznukha Brook, section S-0011, beds 10, 25, 27, Rudny Altai, southwestern Siberia, Russia.



Upper Devonian sections exposed along the right bank of Gryaznukha Brook, near former Razdol'noe Village, SW Rudny Altai and ranges of key radiolarian taxa (for legend see Fig. 3).

Genus *Trilonche* Hinde, 1899, emend. Aitchison & Stratford, 1997, emend. herein

- 1899 *Trilonche* Hinde; Hinde, p. 48.
- 1899 *Staurodruppa* Hinde; Hinde, p. 51.
- 1963 *Entactinosphaera* Foreman; Foreman, p. 274.
- 1988 *Entactinosphaera* Foreman. – Nazarov, p. 60.
- 1999 *Trilonche* Hinde. – Braun & Budil, p. 586.

2000c *Bientactinosphaera* Afanasieva; Afanasieva, pp. 13, 14.

Type species. – *Trilonche vetusta* Hinde, 1899; Upper Devonian, volcanic tuff from Tamworth, New South Wales, Australia.

Original diagnosis. – The test consists of two concentric latticed spheres, with three radial spines at equal or unequal

distances apart. Secondary surface-spines are also sometimes present.

Emended diagnosis. – Further emendment herein after Aitchison & Stratford (1997). Two well-developed spherical or sub-spherical porous or latticed shells and an internal six-rayed double spicule. Six main radial spines developed from rays of the internal spicule. Up to six main spines can be present.

Remarks. – On the basis of thin-section observation of the material, the diagnosis of Hinde (1899, p. 47) restricted the genus to specimens with three radial spines. We agree with Aitchison & Stratford (1997, p. 374) that the genus *Entactinosphaera* reported by Foreman (1963, p. 274) is a junior synonym of *Trilonche* Hinde because the former encompasses material described by Hinde (1899). However, revision of *Entactinosphaera* Foreman made by Nazarov (1975) showed that the mentioned genus is basically composite, including species with two or three shells and a different structure of skeletal sphere. These facts allowed Nazarov to distinguish three different taxa: *Entactinosphaera* Foreman for species with two porous shells; *Spongactinia* Nazarov for species with one spongy and one porous shell; and *Tecentactinia* Nazarov for species with three shells (Nazarov 1975). Thus, Foreman's (1963) diagnosis can be used as a basis for *Trilonche* but it should be corrected by restricting the number of shells strictly to two and including only representatives with porous or latticed structure of the skeletal sphere. Radiolarians with three or more shells and spongy structure should be assigned to other genera.

***Trilonche tanheensis* Luo, Aitchison & Wang, 2002**
Figure 4 K

Holotype. – Figured by Luo, Aitchison & Wang (2002, p. 119, pl. 3, fig. 8), Middle Devonian, Givetian, Tanhe Formation, Wuxiangling Section, Nanning, Guangxi. Collection is deposited in the Department of the Earth Sciences, University of Hong Kong (NKU-WXL128c_14).

Material. – 4 specimens.

Description. – A spherical latticed shell with six three-bladed main spines that are almost equal and robust, disposed at approximately 90° to each other, with 1.25–1.5 torsion, and coiling sinistrally when looking towards the tip. The spines taper rapidly near the tip. By-spines are not developed. The internal structure is not clear.

Dimensions. – Shell diameter 95–105 µm; main spine length 90–110 µm, with base diameter 20–30 µm.

Occurrence. – Upper Devonian, upper Frasnian Gryaznukha Formation; right bank of Gryaznukha Brook, section S-0011, beds 10, 18, 25, 27, SW Rudny Altai, southwestern Siberia, Russia.

Family Haplentactiniidae Nazarov, 1980

Genus *Haplentactinia* Foreman, 1963

Type species. – *Haplentactinia rhinophyusa* Foreman, 1963; Upper Devonian, Frasnian, Huron Member, Ohio, USA.

Diagnosis. – See Foreman (1963, p. 270).

***Haplentactinia rhinophyusa* Foreman, 1963**
Figure 6K, L

1963 *Haplentactinia rhinophyusa* Foreman; Foreman, pl. 1, fig. 2; pl. 3, fig. 7.


2000a *Haplentactinia rhinophyusa* Foreman. – Afanasieva, pl. 25, figs 5, 6.

Holotype. – USNM 640392 figured by Foreman (1963, p. 270, pl. 1, fig. 2), from concretions Norwalk 1, Upper Devonian, Huron Member, Ohio, USA. The holotype and associated collection are deposited in the United States National Museum, Washington D.C.

Material. – 5 specimens.

Description. – The skeletal framework consists of six

Radiolarians from the Gryaznukha Brook sections, Upper Devonian, upper Frasnian, SW Rudny Altai. Scale 50 µm. • A–D – *Cancellactinia acifera* sp. nov., section S-0011; A – holotype, loc. S-0011-25, 1099/1; B – paratype, loc. S-0011-10, 1099/2; C – paratype, loc. S-0011-27, 1099/3; D – paratype, loc. S-0011-10, 1099/4. • E – *Circuliforma robusta* Cheng, section S-0011 (loc. 0011/25), 1099/106. • F – *Palaeoscenidium delicatum* Aitchison, section S-0011 (loc. S-0011-10), 1099/76. • G, H – *Palaeoscenidium cladophorum* Deflandre, section S-0011; G – loc. 00072603/8, 1099/70; H – loc. S-0011-10, 1099/71. • I, J – *Palaeoscenidium tabernaculum* Aitchison, section S-0011 (loc. 00072603/8); I – 1099/77; J – 1099/78. • K, L – *Haplentactinia rhinophyusa* Foreman, section S-0011; K – loc. S-0011-27, 1099/58; L – loc. S-0011-25, 1099/57. • M – *Palaeoscenidium phalagium* Aitchison, section S-0011 (loc. S-0011-25), 1099/79. • N, O – *Ceratoikiscum delicatum* Cheng, section S-0011; N – loc. 00072603/8, 1099/96; O – loc. S-0011-10, 1099/97. • P – *Ceratoikiscum avimexpectans* Deflandre, section S-0011 (loc. 00072603/8), 1099/90. • Q–S – *Nazarovites bioculus* Afanasieva, section S-0011; Q – loc. 00072603/8, 1099/60; R – loc. S-0011-25, 1099/61; S – loc. 00072603/8, 1099/62. • T – *Nazarovites pinnula* Afanasieva, section S-0011 (loc. 00072603/8), 1099/66.



rod-like, slightly tapering spines (rays) that arise from a short medial bar. The internal framework is constructed of one or two groups of branched spinules or apophysis on the spines uniting in a meshwork and creating an isometric latticed shell. The meshwork openings are of angular forms. The distal part of the spines carries a short curved apophysis.

Dimensions. – Shell diameter 100–120 µm; main spine length 90–130 µm, with base diameter 5–10 µm.

Occurrence. – Upper Devonian, upper Frasnian Gryaznukha Formation; right bank of Gryaznukha Brook, section S-0011, beds 18, 25, SW Rudny Altai, southwestern Siberia, Russia.

Family Palacantholithidae Kozur & Mostler, 1981, emend. Afanasieva & Amon, 2005

Genus *Nazarovites* Afanasieva, 2000

Type species. – *Nazarovites bioculus* Afanasieva, 2000; Upper Devonian, middle and upper Frasnian, Domanic Formation, Timan-Pechora Province, northern Russia.

Diagnosis. – See Afanasieva (2000c, p. 4).

Nazarovites bioculus Afanasieva, 2000

Figure 6Q–S

- 1983 *Ceratoikiscum* cf. *spinosiarcuratum* Foreman. – Nazarov & Ormiston, pl. 2, fig. 1.
1997 *Nazarovites bioculus* (nomen nudum); Afanasieva, figs 2–19.
2000a *Nazarovites bioculus* Afanasieva; Afanasieva, pl. 111, figs 1–11.
2000c *Nazarovites bioculus* Afanasieva; Afanasieva, pl. 1, figs 4, 5.

Holotype. – FF-POO1/092-09834 figured by Afanasieva (1997, pp. 7, 8, pl. 1, fig. 5), borehole Shuga-Yag-1003, Middle Frasnian, Domanic Formation, Timan-Pechora Province, Russia. The holotype and associated collection are deposited in the Federal Fund for boring-cores, paleontological and lithological collections, Aprel' Division of All-Russian Research Institute of Petroleum Geology (VNIGNI), Moscow.

Material. – 35 specimens.

Description. – The skeleton is constructed of three main rod-like thin spines intersected in one center at 90°, conditionally vertical (*vL*), horizontal (*gL*) and medial (*mL*). The

apical part of the *vL* spine carries two accessory arch-shaped by-spines (*l*) forming ring structures and situated at the same level. The main and by-spines possess a relatively long, robust, conical apophysis.

Dimensions. – Length of the *mL* spine 280–300 µm; length of the *gL* spine 270–280 µm, with diameter 20 µm; by-spine diameter 17–20 µm; ring structure radius 60 µm; length of the apophysis 10–80 µm, with diameter 5–10 µm.

Occurrence. – Upper Devonian, upper Frasnian Gryaznukha Formation; right bank of Gryaznukha Brook, section S-0011, beds 10, 18, 25, 27, SW Rudny Altai, southwestern Siberia, Russia.

Nazarovites pinnula Afanasieva, 2000

Figure 6T

- 1999 *Ceratoikiscum* sp.; Boundy-Sanders *et al.*, pl. 1, fig. H.
2000a *Nazarovites pinnula* Afanasieva; Afanasieva, pl. 112, figs 1–9.
2000c *Nazarovites pinnula* Afanasieva; Afanasieva, pl. 1, figs 7, 8.

Holotype. – FF-POO1/093-09508 figured by Afanasieva (2000c, pl. 1, fig. 7), Lyayol' River, middle Frasnian, Domanic Formation, Timan-Pechora Province. The holotype and associated collection are deposited in the Federal Fund for boring-cores, paleontological and lithological collections, Aprel' Division of All-Russian Research Institute of Petroleum Geology (VNIGNI), Moscow.

Material. – 35 specimens.

Description. – The skeleton is constructed of three main rod-like thin spines intersected in one center: conditionally, vertical (*vL*) is the most long and thin, horizontal (*gL*) and medial (*mL*) spines are short, and can be thin or robust. Several additional arch-shaped by-spines (*l*) are robust and situated on two levels: the lower spines are slightly curved, and the upper ones are strongly curved forming $\frac{1}{4}$ ring structures. Main and by-spines possess a short, robust conical apophysis.

Dimensions. – Length of the *vL* spine 200–210 µm; length of the *gL* spine 130–150 µm, with diameter 10–12 µm; length of *l* spine 160–210 µm; by-spine diameter 10 µm.

Occurrence. – Upper Devonian, upper Frasnian Gryaznukha Formation; right bank of Gryaznukha Brook, section S-0011, SW Rudny Altai, southwestern Siberia, Russia.

Advanced studies of radiolarian faunas from Upper Devonian strata of the Rudny Altai reveals the domination of spherical entactiniids in the Frasnian. This coincides completely with data known from their global distribution (Nazarov 1988, Braun 1990, Nazarov & Ormiston 1993, Aitchison 1993, Won 1997, Aitchison & Stratford 1997, Aitchison *et al.* 1999, Boundy-Sanders *et al.* 1999, Afanasieva 2000a, Wang *et al.* 2000, Luo *et al.* 2002, Umeda *et al.* 2004). Increasing diversity of ceratoikiscids in the upper Frasnian fauna from the Rudny Altai has already been mentioned in previous reports (Umeda *et al.* 2004, Obut *et al.* 2007). Further detailed study also reveals an abundance of spiny palaeosconids and palacantholithids in the upper Frasnian association.

Continuing radiolarian findings embrace the stratigraphical interval from the top of Givetian to uppermost Frasnian. This data might be used to establish a radiolarian zonation for the Frasnian Stage of SW Siberia.

The authors are grateful to N.V. Sennikov and N.K. Bakharev for guiding us during field studies; E.A. Yolkin for valuable discussions about stratigraphy of the studied area; and I.V. Aborneva for help in taking SEM photographs. The work was supported by grants RFBR No 08-05-00575, SS-3822.2008.5 and from the Russian Science Support Foundation. This is a contribution to UNESCO-IUGS IGCP project 499 “Devonian land-sea interaction: evolution of ecosystems and climate”.

- AFANASIEVA, M.S. 1997. Early Frasnian radiolarians and their biostratigraphic significance. *Doklady Akademii nauk* 355(2), 217–222.
- AFANASIEVA, M.S. 1999. New variant of systematics of Paleozoic radiolarians. *Proceedings of the 13th Geological congress of Komi Republic on geology and mineral resources of the Northeastern European Russia: new results and new prospects*, 253–256. Institut Geologii Komi Nauchnyi Tsentra UB RAS, Syktyvkar. [in Russian]
- AFANASIEVA, M.S. 2000a. *Atlas of Paleozoic Radiolaria of the Russian Platform*. 480 pp. Scientific World, Moscow. [in Russian]
- AFANASIEVA, M.S. 2000b. New radiolarians of Superfamily Entactinoidea from the Upper Devonian of Timan-Pechora Province of Russia. *Paleontologicheskii Zhurnal* 2, 20–33. [in Russian]
- AFANASIEVA, M.S. 2000c. New radiolarians of Orders Aculearia and Sphaerellaria from Upper Devonian of Timan-Pechora Province of Russia. *Paleontologicheskii Zhurnal* 4, 3–19. [in Russian]
- AFANASIEVA, M.S., AMON, E.O., AGARKOV, Y.V. & BOLTOVSKOY, D.S. 2005. Radiolarians in the geological Record. *Paleontological Journal* 39(3), Supplement, 135–392.
- AITCHISON, J.C. 1993. Devonian (Frasnian) radiolarians from the Gogo Formation, Canning Basin, Western Australia. *Palaeontographica, Abteilung A* 228, 105–128.
- AITCHISON, J.C. & STRATFORD, J.M.C. 1997. Middle Devonian (Givetian) Radiolaria from eastern New South Wales, Australia: A reassessment of the Hinde (1899) fauna. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 203, 369–390.
- AITCHISON, J.C., DAVIS, A.M., STRATFORD, J.M.C. & SPILLER, F.C.P. 1999. Lower and Middle Devonian radiolarian biozonation of the Gamilaroi terrane New England Orogen, eastern Australia. *Micropaleontology* 45(2), 138–162. DOI 10.2307/1486110
- BAKHAREV, N.K., SENNIKOV, N.V., YOLKIN, E.A., IZOKH, N.G., ALEKSEENKO, A.A. & OBUT, O.T. 2004. *Key Devonian sections from the Rudny Altai, Salair and Kuznetsk Basin regions*. 104 pp. Publishing house of the Siberian Branch of the Russian Academy of Sciences, Novosibirsk. [in Russian]
- BRAUN, A. 1990. Oberdevonische radiolarien aus Kiesel-schiefer-Geröllen des unteren maintales bei Frankfurt a. M. *Geologischches Jahrbuch Hessen* 118, 5–27.
- BRAUN, A. & BUDIL, P. 1999. A Middle Devonian radiolarian fauna from the Chote Limestone (Eifelian) of the Prague Basin (Barrandian, Czech Republic). In DE WEVER, P. & CAULET, J.-P. (eds) *InterRad VIII, Paris/Bierville 8–13 Septembre 1997. Geodiversitas* 21(4), 581–592.
- BOUNDY-SANDERS, S.Q., SANDBERG, C.A., MURCHEY, B.L. & HARRIS, A.G. 1999. A late Frasnian (Late Devonian) radiolarian, sponge spicule, and conodont fauna from the Slaven Chert, northern Shoshone Range, Roberts Mountains allochthon, Nevada. *Micropaleontology* 45, 62–68. DOI 10.2307/1486203
- CHENG, Y.N. 1986. Taxonomic studies on Upper Paleozoic Radiolaria. *National Museum of Natural Sciences, Taichung, Taiwan, Special Publication* 1, 1–311.
- FOREMAN, H. 1963. Upper Devonian radiolarians from the Huron Member of the Ohio Shale. *Micropaleontology* 9, 267–304. DOI 10.2307/1484751
- HINDE, G.J. 1899. On the Radiolaria in the Devonian rocks of New South Wales. *Quarterly Journal of the Geological Society of London* 55, 38–64.
- LUO, H., AITCHISON, J.C. & WANG, Y.J. 2002. Devonian (upper Emsian–lower Givetian) radiolarians from the Tanhe Formation, Nanning, Guangxi, southwest China. *Micropaleontology* 48, Supplement 1, 113–127.
- LYAKHNITSKY, V.N. & GUTAK, Y.M. 1998. Paleozoic radiolarians from the western part of the Altai-Sayan area (state of art and potential). Geological structure and mineral resources of the western part of the Altai-Sayan Folded Area. *Proceedings of the scientific-practical conference devoted to the 80th anniversary of the West Siberia Geological Survey*, 45–48. Kemerovo-Novokuznetsk. [in Russian]

- NAZAROV, B.B. 1974. Alballlellidae and Paleoscenidiidae from the Upper Devonian beds of the Southern Ural Mountains. *Trudy Vsesoyuznogo Geologicheskogo Instituta, Novaya Seriya* 226, 41–47. [in Russian]
- NAZAROV, B.B. 1975. Lower and Middle Paleozoic radiolarians of Kazakhstan. *Trudy Geologicheskii Institut Akademiia Nauk SSSR* 275, 1–202. [in Russian]
- NAZAROV, B.B. 1980. Discoid Polycystina from the Late Paleozoic, systematics and stratigraphical significance. *Proceedings of the VIII Micropaleontological Meeting, Baku*, 78–79. [in Russian]
- NAZAROV, B.B. 1988. *Paleozoic Radiolarians: Practical manual on microfauna of USSR. Vol. 2.* 232 pp. Nedra, Leningrad. [in Russian]
- NAZAROV, B.B. & ORMISTON, A.R. 1983. Upper Devonian (Frasnian) radiolarian fauna from the Gogo Formation, Western Australia. *Micropaleontology* 29(4), 454–466. DOI 10.2307/1485519
- NAZAROV, B.B. & ORMISTON, A.R. 1993. New biostratigraphically important Paleozoic Radiolaria of Eurasia and North America, 22–60. In BLUEFORD, J.R. & MURCHEY, B. (eds) Radiolaria of giant and subgiant fields in Asia. *Micropaleontology Press, Special Publication 6*.
- OBUT, O.T. 2006. Upper Devonian radiolarians from thin-terrigenous and siliceous strata of the Rudny Altai (South of West Siberia, Russia), 364. In YANG, Q., WANG, Y. & WELDON, E.A. (eds) *Ancient Life and Modern Approaches: Abstracts of the Second International Palaeontological Congress*. University of Science and Technology of China Press.
- OBUT, O.T., IZOKH, N.G. & YOLKIN, E.A. 2007. First occurrences of radiolarians and conodonts in Frasnian siliciclastic sequences of the Rudny Altai (South of West Siberia, Russia). *Subcommission on Devonian Stratigraphy and IGCP 499 Devonian Land Sea Interaction, Eureka, Nevada, 9–17 September 2007*, 69–70. Minuteman Press, Rochester, NY.
- OBUT, O.T., SENNIKOV, N.V. & IWATA, K. 2005. Upper Devonian radiolarians from the Rudny Altai (South of West Siberia). Devonian Terrestrial and Marine Environments: From Continent to Shelf (IGCP 499/SDS joint field meeting). *Contributions of the International Conference. Novosibirsk, July 25–August 9, 2005*, 113–114. Publication House of SB RAS, “Geo Branch”, Novosibirsk.
- RIEDEL, W.R. 1967. Class Actinopoda. Protozoa, 291–298. In HARLAND, W.B., HOLLAND, C.H., HOUSE, M.R., HUGHES, N.F., REYNOLDS, A.B., RUDWICK, M.J.S., SATTERTHWAITTE, G.E., TARLO, L.B.H. & WILLEY, E.C. (eds) *The Fossil Record*. Geological Society of London, London.
- RÜST, D. 1892. Beiträge zur Kenntnis der fossilen Radiolarien aus Gesteinen der Trias und der paläozoischen Schichten. *Palaeontographica, Abteilung A* 38, 107–192.
- SCHWARTZAPFEL, J.A. & HOLDSWORTH, B.K. 1996. Upper Devonian and Mississippian radiolarian zonation and biostratigraphy of the Woodford, Sycamore, Caney and Goddard formations, Oklahoma. *Cushman Foundation for Foraminiferal Research, Special Publication 33*, 1–275.
- UMEDA, M., OBUT, O.T., IWATA, K., SENNIKOV, N.V., IZOKH, N.G. & ERMIKOV, V.D. 2004. Late Devonian radiolarians from the Rudny Altai, Russia. *News of Paleontology and Stratigraphy* 6–7, Supplement to *Geologiya i Geophysica* 45, 103–115.
- WANG, Y.J., FANG, Z.J., YAN, Q., ZHOU, Z.C., CHENG, Y.N., DUAN, Y.X. & XIAO, Y.W. 2000. Middle-Late Devonian strata of cherty facies and radiolarian faunas from west Yunnan. *Acta Micropalaeontologica Sinica* 17, 235–254.
- WEDDIGE, K. Ed. 2006. Devonian Correlation Table. Supplements 2006. *Senckenbergiana Lethaea* 86(2), 177–197.
- WON, M.-Z. 1997. Review of family Entactiniidae (Radiolaria), taxonomy and morphology of Entactiniidae in the Late Devonian (Frasnian) Gogo Formation, Australia. *Micropaleontology* 43(4), 333–369. DOI 10.2307/1485930
- YOLKIN, E.A., BAKHAREV, N.K., IZOKH, N.G., GRATSIAKOVA, R.T., KIPRIYANOVA, T.P. & OBUT, O.T. 2005. Devonian sequences of Salair, Rudny and Gorny Altai. Field Excursion Guidebook. *International Conference “Devonian Terrestrial and Marine Environments: from Continent to Shelf” (IGCP 499 Project / SDS joint field meeting). Novosibirsk, Russia. July 25–August 9, 2005*. 80 pp. Publishing House of SB RAS, “Geo” Branch, Novosibirsk.
- YOLKIN, E.A., KONTOROVICH, A.E. & BAKHAREV, N.K. 2003. Facial megazones in the Paleozoic of the West Siberia Geosyncline. *Geodynamic evolution of the lithosphere of the Central-Asian Folded Belt. Contributions to the scientific session on Basic Research Program (20–23 October, 2003, Irkutsk)*, 100–104. Irkutsk. [in Russian]
- YOLKIN, E.A., IZOKH, N.G. & BAKHAREV, N.K. 2006. Annotations to the Devonian Correlation Tables, R 175–184, H 110–111, R 608–612 di-ds 06: Devonian of West Siberia (Salair, Kuznetsk Basin, Gorny [High] and Rudny [Ore] Altai). *Senckenbergiana Lethaea* 86(2), 173–175.