

11. References

- Anderson D.E. & Olimpio J.C. (1977): Progressive homogenization of metamorphic garnets, South Moray, Scotland: Evidence for volume diffusion.* - *Canad. Mineral.*, 15, 205-216. Ottawa.
- Anovitz L.M. (1991): Al zoning in pyroxene and plagioclase: Window on late prograde to early retrograde P-T paths in granulite terranes.* - *Amer. Mineral.*, 76, 1328-1343. Washington.
- Behr H.-J. (1978): Subfluenz-Prozesse im Grundgebirge Stockwerk Mitteleuropas.* - *Z. Dtsch. geol. Gesell.*, 129, 283-318. Hannover.
- Behr H.-J., Fritsch E. & Mannsfeld L. (1965): Die Granulite von Zöblitz im Erzgebirge als Beispiel für Granulitbildung in tiefreichenden Scherhorizonten.* - *Krystalinikum*, 3, 7-29. Praha.
- Behr H.-J., Engel W. & Franke W. (1982): Variscan wildflysch and nappe tectonics in the Saxothuringian zone (northeast Bavaria, West Germany).* - *Amer. J. Sci.*, 282, 1438-1470. New Haven.
- Bell T.H., Fleming P.D. & Rubenach M.J. (1986): Porphyroblast nucleation, growth and dissolution in regional metamorphic rocks as a function of deformation partitioning during foliation development.* - *J. metamorph. Geol.*, 4, 37-67. Oxford.
- Bence A.E. & Albee A.L. (1968): Empirical correction factors of the electron microanalyses of silicates and oxides.* - *J. Geol.*, 76, 382-403. Chicago.
- Benoit V. & Mercier J.-C. (1986): Les enclaves ultramafiques du volcanisme alcalin tertiaire du centre du plateau du Colorado: implications tectoniques.* - *Bull. Soc. géol. France*, 8, 1015-1023. Paris.
- Boettcher A.L. (1970): The system CaO-Al₂O₃-SiO₂-H₂O at high pressures and temperatures.* - *J. Petrol.*, 11, 337-379. Oxford.
- Bohlen S.R. (1987): Pressure-temperature-time paths and a tectonic model for the evolution of granulites.* - *J. Geol.*, 95, 617-632. Chicago.
- Boland J.N. & van Roermund H.L.M. (1983): Mechanism of exsolution in omphacites from high-temperature, type B, eclogites.* - *Phys. Chem. Minerals*, 9, 30-37. Berlin.
- Brey G.P. & Köhler T. (1990): Geothermobarometry in four-phase Iherzolites II. New thermobarometers, and practical assessment of existing thermobarometers.* - *J. Petrol.*, 31, 1353-1378. Oxford.
- Carswell D.A. (1991): Variscan high P-T metamorphism and uplift history in the Moldanubian Zone of the Bohemian Massif in Lower Austria.* - *Eur. J. Mineral.*, 3, 323-342. Stuttgart.
- Carswell D.A. & Cuthbert S.J. (1986): Eclogite facies metamorphism in the lower continental crust.* - In J.B. Dawson, D.A. Carswell, J. Hall and K.H. Wedepohl (eds), *The Nature of the Lower Continental Crust.* - *Geol. Soc. Spec. Publ.* 24, 193-209.
- Carswell D.A. & Gibb F.G.F. (1987): Evaluation of mineral thermometers and barometers applicable to garnet Iher-*
- zolite assemblages.* - *Contr. Mineral. Petrol.*, 95, 499-511. Berlin - Heidelberg - New York.
- Chaloupský J. (1977): Stratigraphie du Precambrien du Massif de Bohême. In La chaîne Varisque d'Europe moyenne et occidentale. Coll. intern. CNRS, 243, 17-32. Rennes.*
- Chaloupský J. (1989): Major tectonostratigraphic units of the Bohemian Massif.* - *Geol. Soc. Am. Spec. Paper*, 230, 101-114. New York.
- Chatterjee N.D. (1976): Margarite stability and compatibility relations in system CaO-Al₂O₃-SiO₂-H₂O.* - *Amer. Mineral.*, 60, 985-993. Washington.
- Clemens J.D. (1990): The granulite - granite connection.* - In D. Vielzeuf and Ph. Vidal (eds), *Granulites and Crustal Evolution.* - Kluwer Academic Publishers, Netherlands.
- Cressey G., Schmid R. & Wood B.J. (1978): Thermodynamic properties of almandine-grossular garnet solid solutions.* - *Contr. Mineral. Petrol.*, 67, 397-404. Berlin - Heidelberg - New York.
- Cullers R.L. & Graf J.L. (1984): Rare earth elements in igneous rocks of the continental crust: intermediate and silicic rocks - ore petrogenesis.* In P. Henderson (ed.), *Rare earth element geochemistry*, 275-316. Elsevier, Amsterdam.
- Dathe E. (1882): Beiträge zur Kenntnis des Granulits.* - *Z. Dtsch. geol. Gesell.*, 34, 12-40. Hannover.
- Droop G.T.R. & Bucher-Nurminen K. (1984): Reaction textures and metamorphic evolution of sapphirine-bearing granulites from the Gruf Complex, Italian central Alps.* - *J. Petrol.*, 25, 766-803. Oxford.
- Ebert H. (1932): Der Varistische Untergrund Nordböhmens zwischen Erzgebirge und Jeschken.* - *Z. Dtsch. geol. Gesell.*, 84, Hannover.
- El Bouseily A.M. & El Sokkary A.A. (1975): The relation between Rb, Ba, and Sr in granite rocks.* - *Chem. Geol.*, 16, 207-219. Amsterdam.
- Ellis D.J. (1987): Origin and evolution of granulites in normal and thickened crusts.* - *Geology*, 15, 167-170. Boulder.
- Ellis D.J. & Green E.H. (1979): An experimental study of the effect of Ca upon garnet-clinopyroxene Fe-Mg exchange equilibria.* - *Contr. Mineral. Petrol.*, 71, 13-22. Berlin - Heidelberg - New York.
- England P.C. (1987): Diffuse continental deformation: length scales, rates and metamorphic evolution.* In E.R. Oxburgh, B.W.D. Yardley and P.C. England (eds), *Tectonic Settings of Regional Metamorphism*, 3-22. The Royal Society, Cambridge University Press, London.
- England P.C. & Thompson A.B. (1984): Pressure-temperature paths of regional metamorphism. I. Heat transfer during the evolution of regions of thickened continental crust.* - *J. Petrol.*, 25, 894-928. Oxford.
- England P.C. & Thompson A.B. (1986): Some thermal and tectonic models for crustal melting in continental collision zones.* In M.P. Coward and A.C. Ries (eds.), *Collision tectonics.* - *Geol. Soc. Lond., Spec. Publ.* No. 19, 83-94. London.
- Fediuková E. (1978): Mafic minerals from granulites of the borehole Holubov (South Bohemian Moldanubicum).* - *Sbor. geol. Věd, Lož. Geol. - Mineral.*, 19, 169-198, Praha.
- Ferry J.M. & Spear F.S. (1978): Experimental calibration of*

- the partitioning of Fe and Mg between biotite and garnet. - *Contr. Mineral. Petrol.*, 66, 113-117. Berlin - Heidelberg - New York.
- Fiala J. (1965): Pyrope of some garnet peridotites of the Czech Massif. - Krystalinikum*, 3, 55-74, Praha.
- Fiala J., Matějovská O. & Vaříková V. (1987): Moldanubian granulites and related rocks: petrology, geochemistry and radioactivity. - Rozpr. Čs. Akad. Věd, Ř.mat.přír.Věd*, 97, 102pp., Academia Praha.
- Fiala J. & Paděra K. (1977): Eclogitic rocks from the basement of the České Středohoří Mts. - Krystalinikum*, 17, 33-45. Praha.
- Fiala J. & Paděra K. (1984): The chemistry of the minerals of the pyrope dunite from borehole T-7 near Staré (Bohemia). - Tschermaks Min. Petr. Mitt., TMPM*, 24, 205-219. Wien.
- Fiala J. & Vaříková V. (1985): Radioactivity and geochemistry of granulites and some related rocks from the Saxonian Granulite Complex. - Stud. geophys. geod.*, 29, 374-394. Academia Praha.
- Franke W. (1989): Tectonostratigraphic units in the Variscan belt of central Europe. - Geol. Soc. Am. Spec. Paper*, 230, 67-90. New York.
- Frost B.R. & Chacko T. (1989): The granulite uncertainty principle: limitations on geothermobarometry in granulites. - J. Geol.*, 97, 435-450. Chicago.
- Frost B.R., Frost C.D. & Touret J.R.L. (1989): Magmas as a source of heat and fluids in granulite metamorphism. - In D.Bridgwater (ed.), Fluid Movements - Element Transport and the Composition of the Deep Crust, I-18. - Kluwer Academic Publishers, Netherland.*
- Fuhman M.L. & Lindsley D.H. (1988): Ternary-feldspar modelling and thermometry. - Am. Mineral.*, 73, 201-215. Washington.
- Fyfe W. (1973): The granulite facies, partial melting and the Archean crust. - Phil. Trans. Roy. Soc. Lond.*, A273, 457-461. London.
- Gardien V., Tegyey M., Lardeaux J.-M., Misseri M. & Dufour E. (1990): Crust-mantle relationships in the French Variscan Chain: the example of the Southern Monts du Lyonnais unit (eastern French Massif Central). - J. metamorph. Geol.*, 8, 477-492. Oxford.
- Gasparik T. & Lindsley D.H. (1980): Phase equilibria at high pressure of pyroxenes containing monovalent and trivalent ions. In C.T.Prewitt (ed.), Pyroxenes. - Mineral. Soc. Am. Rev. Mineral.*, 7, 309-339. Washington.
- Gebauer D. (1991): Two Palaeozoic high-pressure events in a garnet-peridotite of northern Bohemia, Czechoslovakia. - Abstract. 2nd Eclogite Field Symp., Granada. Terra Abstracts. Terra Nova 3, Oxford.*
- Goldsmith J.R. (1980): The melting and breakdown reactions of anorthite at high pressures and temperatures. - Amer. Mineral.*, 65, 272-284. Washington.
- Gray C.M. (1977): The geochemistry of Central Australian granulites in relation to the chemical and isotopic effects on granulite facies metamorphism. - Contr. Mineral. Petrol.*, 65, 79-89. Berlin - Heidelberg - New York.
- Green D.H. & Ringwood A.E. (1967): An experimental investigation of gabbro to eclogite transition and its petrological applications. - Geochim. cosmochim. Acta*, 31, 767-833. Oxford.
- Green T.H., Brunfelt A.O. & Heier K.S. (1972): Rare-earth element distribution and K/Rb ratios in granulites, mangerites and anorthosites, Lofoten-Vesteraalen, Norway. - Geochim. cosmochim. Acta*, 36, 241-257. Oxford.
- Guidotti Ch. V. (1984): Micas in metamorphic rocks. In S.W.Bailey (ed), Micas. Mineral. Soc. Am. Rev. Mineral.*, 13, 357-456. Washington.
- Harley S.L. (1984): An experimental study of the partitioning of Fe and Mg between garnet and orthopyroxene. - Contr. Mineral. Petrol.*, 86, 359-373. Berlin - Heidelberg - New York.
- Harley S.L. (1989): The origins of granulites: a metamorphic perspective. - Geol. Mag.*, 126, 215-247. Cambridge.
- Haskin L.A. (1984): Petrogenetic modelling - use of rare earth elements. In P. Henderson (ed.), Rare earth element geochemistry, 115-152. Elsevier. Amsterdam.*
- Haskin L.A., Frey F.A., Schmitt R.A. & Smith R.H. (1966): Meteoric, solar and terrestrial rare-earth distributions. - Phys. Chem. Earth*, 7, 169-321. London.
- Henderson P. (ed.) (1984): Rare earth element geochemistry. - Developments in Geochemistry*, 2., 510p. - Elsevier, Amsterdam.
- Hey M.H. (1954): A new review of the chlorites. - Min. Mag.*, 30, 277. London.
- Hibsch J.E. (1920): Geologische Karte des Böhmischen Mittelgebirges, Blat. XIV. (Meronitz - Trebnitz)*
- Hodges K.V. & Spear F.S. (1982): Geothermometry, geobarometry and the Al_2SiO_5 triple point at Mt. Moosilauke, New Hampshire. - Amer. Mineral.*, 67, 1118-1134. Washington.
- Hokr Z. et al. (1974): Základní geologická mapa 1:25 000, list M 33-51-D-a Prunéřov. Ústř. Úst. Geol.*, Praha.
- Holdaway M.J. (1981): Stability of andalusite and the aluminum silicate phase diagram. - Amer. J. Sci.*, 271, 97-131. New Haven.
- Holdaway M.J. & Lee S.M. (1977): Fe-Mg cordierite stability in high grade pelitic rocks based on experimental and natural observations. - Contr. Mineral. petrol.*, 63, 175-198. Berlin - Heidelberg - New York.
- Holland T.J.B. (1979): Reversed hydrothermal determination of jadeite-diopside activities. - EOS*, 60, 405.
- Holland T.J.B. (1980): The reaction albite = jadeite + quartz determined experimentally in the range 600-1200°C. - Amer. Mineral.*, 65, 129-134. Washington.
- Holubec J. (1956): Geologické-petrografické poměry krysalinika v okoli Místa (záp. od Chomutova) v Krušných horách. - Sbor. Geol. Úst. ČSR*, 23, 111-158, Praha.
- Hsu L.C. (1968): Selected phase relationships in the system Al-Mn-Fe-Si-O; a model for garnet equilibria. - J.Petrol.*, 9, 40-43. Oxford.
- Humphris S.E. (1984): The mobility of the rare earth elements in the crust. In P. Henderson (ed.), Rare earth element geochemistry, 317-342. Elsevier. Amsterdam.*
- Hunt J.A. & Kerrick D.M. (1977): The stability of sphene: Experimental redetermination and geological implications. - Geochim. cosmochim. Acta*, 41, 279-288. Oxford.

- Ibrmajer J. & Suk M. et al. (1989): Geofyzikální obraz ČSSR. - Academia, Praha.
- Indares A. & Martignole J. (1985a): Biotite-garnet geothermometry in granulite facies rocks: evaluation of equilibrium criteria. - Canad. Mineral., 23, 187-193. Ottawa.
- Indares A. & Martignole J. (1985b): Biotite-garnet geothermometry in the granulite facies: the influence of Ti and Al in biotite. - Amer. Mineral., 70, 272-278. Washington.
- Jakeš P. (1969): Retrogressive changes of granulite-facies rocks - an example from the Bohemian Massif. - Spec. Publ. geol. Soc. Aust., 2, 367-374. Canberra.
- Jakeš P. & Taylor S.R. (1974): Excess europium content in Precambrian sedimentary rocks and continental evolution. - Geochim. cosmochim. Acta, 38, 739-745. Oxford.
- Jakeš P., Jelinek E. & Misař Z. (1985): Ultramafic rocks within granulites - a type of crust-mantle transition in mountain belts. - Abstract. EUG III, Strasbourg.
- James R.S. & Hamilton D.L. (1969): Phase relations in the system NaAlSi₃O₈ - KAlSi₃O₈ - CaAl₂Si₂O₈-SiO₂ at 1 kilobar water vapour pressure. - Contr. Mineral. Petrol., 21, 111-141. Berlin - Heidelberg - New York.
- Jenkins D.M. (1981): Experimental phase relations of hydrous peridotites modelled in the system H₂O-CaO-MgO-Al₂O₃-SiO₂. - Contr. Mineral. Petrol., 77, 166-176. Berlin - Heidelberg - New York.
- Jensen L.S. (1976): A new cation plot for classifying subalkalic volcanic rocks. - Ontario Div. Mines, MP 66, 22p. Toronto.
- Joanny V., van Roermund H. & Lardeaux J.M. (1991): The clinopyroxene/plagioclase symplectite in retrograde eclogites: a potential geothermobarometer. - Geol. Rndsch., 80, 303-320. Stuttgart.
- Klárová H. (1990): Eclogites of the Bohemian Part of the Saxothuringicum. - Rozpr. Čs Akad. Věd, Ř.mat.přír.Věd, 100, 5, 1-86. Praha.
- Klárová H., Lardeaux J.-M. & Schulmann K. (1992): Tectono-metamorphic evolution of the Saxothuringian eclogites (Central Erzgebirge): A record of superposed nappe tectonics. - Abstract. 7th geological workshop Styles of Superposed Variscan Tectonics, Kutná Hora. ČSAV Praha.
- Kodym O., Jakeš P. & Schovánek P. (1978): Granulite und ultramafische Gesteine aus der Strukturborung Holubov. - Sbor. geol. Věd, Geologie, 32, 7-48. Praha.
- Koops P.O & Thompson A.B. (1985): Non-mafic rocks in the greenschist, blueschist and eclogite facies. - Chem. Geol., 50, 3-30. Amsterdam.
- Kopecký L. (1975): Geologické mapování na listu Kadaň, M 33-51-D-c. - Archiv Ústř.úst. Geol., Praha.
- Kopecký L. (1986): Geologic development and block structure of the Cenozoic Ohře rift (Czechoslovakia). In M.J. Aldrich and A.W. Laughlin (eds), Proc. 6th Internat. Conf. Basement Tectonics, Los Alamos, New Mexico, USA, 114-124. Los Alamos.
- Kopecký L. (1987, 1988): Mladý vulkanismus Českého masivu. - Geologie a hydrometallurgie uranu, 11 & 12. Stráž pod Ralskem.
- Kopecký L. & Paděra K. (1974): Bänderung der Ultramafitischen Gesteine in der Bohrung T-7-Staré bei Třebenice (Nordböhmien). - In E. Alexiev, J. Minčeva-Stefanova and T.G. Radonova (eds), Minerogenesis, Bulgar. Acad. Sci. Geol. Inst., 161-169. Sofia.
- Kopecký L. & Satran V. (1966): Buried occurrences of pyrope-peridotite and the structure of the crystalline basement in the extreme SW of the České středohoří mountains. - Krystalinikum, 4, 65-86. Praha.
- Kopecký L., Píšová J., Ložek V. et al. (1960): Mineralogicko-petrografický výzkum pyroponosných ultrabasik Českého středohoří. Zpráva za rok 1958 a 1959. - MS, Ústř. úst. Geol., Praha.
- Kopecký L., Píšová J., Satran V & Pokorný V. (1961): Mineralogicko-petrografický výzkum pyroponosných ultrabasik Českého středohoří. Mineralogický, petrografický a geofyzikální výzkum v oblasti Českého středohoří a v severových. Čechách. MS, Ústř. úst. Geol., Praha.
- Kopecký L., Dobeš M., Fiala J. & Štovičková N. (1970): Fenites of the Bohemian Massif and the relations between fenitization, alkaline volcanism and deep fault tectonics. - Sborník geol. Věd, Geologie, 16, 51-112. Academia, Praha.
- Kotková J. (1991): Geotermobarometrie a trajektorie P-T-t v horninách granulitové facie. - MS, práce ke kandid. minimu, 120p., Ústř. úst. Geol., Praha.
- Kotková J. (1992): Granulites and associated peridotites of north Bohemia (Bohemian Massif). - PhD Thesis. MS, Czech Geological Survey, Praha.
- Kozlo A.M. & Newton R.C. (1988): Redetermination of the anorthite breakdown reaction and improvement of the plagioclase-garnet-Al₂SiO₅-quartz geobarometer. - Amer. Mineral., 73, 216-223. Washington.
- Kozlo A.M. & Newton R.C. (1989): Grossular activity-composition relationships in ternary garnets determined by reversed displaced-equilibrium experiments. - Contr. Mineral. Petrol., 103, 423-433. Berlin - Heidelberg - New York.
- Köhler T.P. & Brey G.P. (1990): Calcium exchange between olivine and clinopyroxene calibrated as a geotermobarometer for natural peridotites from 2 to 60 kb with applications. - Geochim. cosmochim. Acta., 54, 2375-2388. Oxford.
- Kröner A., Frischbutter A., Bergner R. & Hofmann J. (1991): Zirkon-Evaporationsalter von granitoiden Gesteinen aus dem Erzgebirge und dem Rande der Lausitz und ihre geodynamische Bedeutung. - Kurzfasungen 7. Rundgespräch Geodynamik des europäischen Varisziums, Freiberg.
- Kushiro I. (1969): Clinopyroxene solid solutions formed by reaction between diopside and plagioclase at high pressure. In 'Pyroxenes and amphiboles: crystal chemistry and phase petrology'. - Contr. Mineral. Petrol., 66, 75-80. Berlin - Heidelberg - New York.
- Lambert I.B. & Heier K.S. (1967): The vertical distribution of uranium, thorium, and potassium in the continental crust. - Geochim. cosmochim. Acta, 31, 377-390. Oxford.
- Lambert I.B. & Heier K.S. (1968): Geochemical investigations of deep-seated rocks in the Australian shield. - Lithos,

- 1, 30-53. Oslo.
- Lardeaux J.-M., Nisio P. & Boudeulle M. (1987):* Deformational and metamorphic history at the Lago Superiore area of the Monviso ophiolitic complex (Italian western Alps): A record of subduction-collision cycle? - *Ophioliti*, 12, 479-502. Bologna.
- Lardeaux J.-M., Caron J.-M., Nisio P., Péquignot G. & Boudeulle M. (1986):* Microstructural criteria for reliable thermometry in low-temperature eclogites. - *Lithos*, 19, 187-203. Oslo.
- Leake B.E. (1978):* Nomenclature of amphiboles. - *Amer. Mineral.*, 63, 1023-1052. Washington.
- LeBreton N. & Thompson A.B. (1988):* Fluid-absent (dehydration) melting of biotite in metapelites in the early stage of crustal anatexis. - *Contr. Mineral. Petrol.*, 99, 226-237. Berlin - Heidelberg - New York.
- Ledru P., Lardeaux J.-M., Santallier D., Autran A., Quenardel J.-M., Floc'h J.-P., Lerouge G., Maillet N., Marchand J. & Ploquin A. (1989):* Où sont les nappes dans le Massif central français? - *Bull. Soc. géol. France*, 8, 605-618. Paris.
- LeGoff E. & Ballévre M. (1990a):* Geothermobarometry in albite-garnet orthogneisses: A case study from the Gran Paradiso Nappe (Western Alps). - *Lithos*, 25, 261-280. Oslo.
- LeGoff E. & Ballévre M. (1990b):* Méthodes d'estimation des conditions pression-température dans les orthogneiss: analyse des relations de phases. - *C.R. Acad. Sci. Paris*, t.311, II, 119-125. Paris.
- Libourel G. (1985):* Le complexe de Santa Lucia di Mercurio (Corse). Ultramafites mantelliques, intrusion basique stratifié, paragneiss granulitiques. Un équivalent possible des complexes de la zone d'Ivrée. - Thèse 3ème cycle. Université P.Sabatier, Toulouse.
- Massone J.J. & Schreyer W. (1987):* Phengite barometry based on the limiting assemblage with K-feldspar, phlogopite and quartz. - *Contr. Mineral. Petrol.*, 96, 212-224. Berlin - Heidelberg - New York.
- Matějkovská O. (1967):* Petrogenesis of the Moldanubian granulites near Náměšť nad Oslavou. - *Krystalinikum*, 5, 85-103. Praha.
- Matějkovská O & Vařková V. (1980):* The concentrations of radioactive and trace elements in the granulites from the south-east margin of the Bohemian Massif (Czechoslovakia). - *Stud. geophys. geod.*, 24, 285-301. Academia, Praha.
- McLennan S.M. & Taylor S.R. (1980):* Th and U in sedimentary rocks: Crustal evolution and sedimentary recycling. - *Nature*, 285, 621-624. London.
- Medaris L.G. & Carswell D.A. (1990):* The petrogenesis of Mg-Cr garnet peridotites in European metamorphic belts. - In D.A.Carswell (ed), *Eclogite Facies Rocks*. - Blackie.
- Mercier L., van Roermund H.L.M. & Lardeaux J.M. (1991):* Comparison of PT paths in allochthonous high pressure metamorphic terrains from the Scandinavian Caledonides and the French Massif Central: contrasted thermal structures during uplift. - *Geol. Rndsch.*, 80, 333-348. Stuttgart.
- Mezger K. (1990):* Geochronology in granulites. - In D.Vielzeuf and Ph.Vidal (eds), *Granulites and Crustal Evolution*. - Kluwer Academic Publishers, Netherlands.
- Mineyev D.A. (1963):* Geochimickaja differenciacija redkozemelnich elementov (Geochemical differentiation of the rare-earths). - *Geochimija*, 12, 1129-1149. Moskva.
- Misař Z., Jelínek E. & Jakeš P. (1984):* Inclusions of peridotite, pyroxenite and eclogite in granulite rocks of pre-Hercynian mantle and lower crust in the eastern Bohemian Massif (Czechoslovakia). - *Ann. Sci. Univ. Clermont Fd*, II, 74, 85-95. Clermont-Ferrand.
- Milčoch B. (1984):* Základní geologická mapa ČSSR 1:25 000, list 02-314 Litvínov. - MS, Ústř.úst. Geol., Praha.
- Milčoch B. & Schulmann K. (1992):* Superposition of Variscan ductile shear deformation on pre-Variscan mantled gneiss structure (Catherine dome, Erzgebirge, Bohemian Massif). - *Geol. Rndsch.*, 81/2, 501-513. Stuttgart.
- Mori T. & Green D.H. (1978):* Laboratory duplication of phase equilibria observed in natural garnet Iherzolites. - *J. Geol.*, 86, 83-97. Chicago.
- Möbus G. (1964):* Die geotektonische entwicklung des grundgebirges im raum Erzgebirge - Elbtalzone - Lausitzer Grundgebirge - Westsudeten. - Abh. Deutsch. Akad. Wiss. Berlin, K. Chemie, Geol., Biol., 5, 5-114. Akademie Verlag, Berlin.
- Newton R.C. (1966):* Some calc-silicate equilibrium relations. - *Amer. J. Sci.*, 264, 204-222. New Haven.
- Newton R.C. (1989):* Metamorphic fluids in the deep crust. - *Ann. Rev. Earth Planet. Sci.*, 17, 385-412. Amsterdam.
- Newton R.C. & Haselton H.T. (1981):* Thermodynamics of the garnet-plagioclase-Al₂SiO₅-quartz geobarometer. In: Newton et al. (eds), *Thermodynamics of Minerals and Melts*, 131-147. Springer-Verlag, New York.
- Newton R.C., Charlu T.V. & Kleppa O.J. (1980a):* Thermochemistry of the high structural state plagioclase. - *Geochim. cosmochim. Acta*, 44, 933-941. Oxford.
- Newton R.C., Smith J.V. & Windley B.F. (1980b):* Carbonic metamorphism, granulites and crustal growth. - *Nature*, 288, 45-50. London.
- Nickel K.G. & Green D.H. (1985):* Empirical geothermobarometry for garnet peridotites and implications for the nature of the lithosphere, kimberlites and diamonds. - *Earth Planet. Sci. Lett.*, 73, 158-170. Amsterdam.
- Obata M. & Morten L. (1985):* The spinel Iherzolite to garnet Iherzolite transformation and the gravity sinking of ultramafic bodies in the lower crust. - *Terra Cognita* 5, 440. Wien.
- O'Hara M.J., Richardson S.W. & Wilson G. (1971):* Garnet peridotite stability and occurrence in crust and mantle. - *Contr. Mineral. Petrol.*, 32, 48-68. Berlin - Heidelberg - New York.
- O'Neill H.S.C. (1981):* The transition between spinel Iherzolite and garnet Iherzolite, and its use as a geobarometer. - *Contr. Mineral. Petrol.*, 77, 185-194. Berlin - Heidelberg - New York.
- O'Neill H.S.C. & Wood B.J. (1979):* An experimental study of Fe-Mg partitioning between garnet and olivine and its calibration as a geothermometer. - *Contr. Mineral. Petrol.*, 70, 59-70. Berlin - Heidelberg - New York.
- Opletal M. & Vrána S. (1989):* Xenolithy chamoxykitových

- hornin ve vulkanitech Českého středohoří. - Sbor. geol. Věd, Geologie, 44, 51-78. Praha.
- Oxburgh E.R. & Turcotte D.L. (1974): Thermal gradients and regional metamorphism in overthrust terrains with special reference to the Eastern Alps.* - Schweiz. mineral. petrogr. Mitt., 54, 641-662. Zürich.
- Pecerillo A. & Taylor R.S. (1976): Geochemistry of Eocene calc-alkaline volcanic rocks from Kastamonu area, Northern Turkey.* - Contr. Mineral. Petrol., 68, 63-81. Berlin - Heidelberg - New York.
- Perkins D., Westrum E.F. & Essene E.J. (1980): Thermodynamic properties and phase relations of some minerals in the system CaO - Al₂O₃ - SiO₂ - H₂O.* - Geochim. cosmochim. Acta, 44, 61-84. Oxford.
- Petrakakis K. (1986): Metamorphism of high-grade gneisses from the Moldanubian zone, Austria, with particular reference to the garnets.* - J. metamorph. Geol., 4, 323-344. Oxford.
- Pin Ch. & Pecaut J.-J. (1986): Ages des épisodes de métamorphisme paléozoïques dans le Massif central et le Massif armoricain.* - Bull. Soc. géol. France, 8, 461-469. Paris.
- Pin Ch. & Vielzeuf D. (1983): Granulites and related rocks in Variscan median Europe: A dualistic interpretation.* - Tectonophysics, 93, 47-74. Amsterdam.
- Pin Ch. & Vielzeuf D. (1988): Les granulites de haute-pression d'Europe moyenne témoins d'une subduction éo-hercynienne. Implications sur l'origine des groupes leptyno-amphiboliques.* - Bull. Soc. géol. France, 8, 13-20. Paris.
- Pokorný V. & Štovičková N. (1980): Geofyzikální projevy strukturně geologických deformací v Českém masivu.* - MS, Geofyzika Praha.
- Pouba Z., Fiala J. & Paděra K. (1985): Omphacite granulite from the NE marginal area of the Bohemian Massif (Rychleby Mts.).* - Neu. Jb. Mineral., Abh., 151, 29-52. Stuttgart.
- Powell R. (1985): Regression diagnostics and robust regression in geothermometer/geobarometer calibration: the garnet-clinopyroxene geothermometer revisited.* - J. metamorph. Geol., 3, 231-243. Oxford.
- Powell R. & Holland T.J.B. (1988): An internally consistent dataset with uncertainties and correlations: 3. Applications to geobarometry, worked examples and a computer program.* - J. metamorph. Geol., 6, 173-204. Oxford.
- Pride C. & Muecke G.K. (1980): Rare earth geochemistry of the Scourian Complex N.W.Scotland - evidence for the granite-granulite link.* - Contr. Mineral. Petrol., 73, 403-412.
- Ramsay J.G. (1980): Shear zone geometry: a review.* - J. struct. Geol., 2, 83-100. Bristol.
- Ramsay J.G. & Graham (1970): Strain variation in shear belts.* - Can. J. Earth Sci., 7, 786-813. Ottawa.
- Rogers J.J.W. & Adams J.A.S. (1978): Th: Abundances in common igneous rocks.* In K.H. Wedepohl (ed), Handbook of Geochemistry, 90-E-1 - 90-E-12. Springer-Verlag, Berlin.
- Rost M. (1928): Geologie des crusallinen Grundgebirges am Erzgebirgsrand zwischen Keilberg und Klösterle.* - Abh. Sächs. geol. Landesamts, 7, Leipzig.
- Rost F. & Grigel W. (1959): Zur Geochemie und Genese granatführender Ultramafitite des mitteleuropäischen Grundgebirges.* - Chem. d. Erde, 28, 91-177. Jena.
- Rötzler J. (1992): Zur Petrogenese im Sächsischen Granulitgebirge. Die pyroxenfreien Granulite und die Metapelite.* - Geotek. Forsch., 77, 1-100. Stuttgart.
- Rötzler K. (1992): High-pressure metamorphism of quartz-feldspathic granulites in the Erzgebirge, southern Saxony, Germany.* - Abstract. International workshop High pressure granulites - lower crustal metamorphism, Rohanov. ČGÚ Praha.
- Rubie D.C. & Thompson A.B. (1985): Kinetics of metamorphic reactions at elevated temperatures and pressures: an appraisal of available experimental data.* In A.B. Thompson and D.C. Rubie (Eds), Metamorphic reactions, kinetics, textures, and deformation. Springer-Verlag, New York.
- Rudnick R.L. & Presper T. (1990): Geochemistry of intermediate- to high-pressure granulites.* - In D. Vielzeuf and Ph. Vidal (eds), Granulites and Crustal Evolution. - Kluwer Academic Publishers, Netherlands.
- Rudnick R.L., McLennan S.M. & Taylor S.R. (1985): Large ion lithophile elements in rocks from high-pressure granulite facies terrains.* - Geochim. cosmochim. Acta, 49, 1345-1655. Oxford.
- Sattran V. (1964): Geologické mapování krystalinika na listu 1:50 000 Chomutov. M33-51-D-a (Prunéřov).* MS Ústř. úst. Geol., Praha.
- Sattran V. (1966): Zpráva o základním geologickém výzkumu krušnohorského krystalinika mezi Mistem a Chomutovem.* - Zprávy o geol. výzkumech v r. 1964. Praha.
- Sattran V. (1967): Krystalinikum údolí Ohře.* In Kopecký L. et al.: Vysvětlivky k listu mapy 1:50 000 M33 51 C(c,d). MS, Ústř. úst. Geol., Praha.
- Sattran V. & Kopecký L. (1967): Podklad pro mapu 1:50 000 list Vejprty. M33-51-C(c,d)(Perštějn, Kadaň).* - MS, Ústř. úst. Geol., Praha.
- Scharbert H.G., Korkisch J. & Stefan I. (1976): Uranium, thorium and potassium in granulite facies rocks, Bohemian Massif, Lower Austria, Austria.* - Tschermaks Min. Petr. Mitt. TMPM, 23, 223-232. Wien.
- Schmädicke E., Schmidt W. & Okrusch M. (1991): Zur polymetamorphen Entwicklung der Eklogite des Erzgebirges.* - Kurzfassungen 7. Rundgespräch Geodynamik des europäischen Variszikums. Freiberg.
- Šrbený O. & Ullrich J. (1988): Crystalline basement of the České středohoří Mountains as inferred from the inclusions in Tertiary volcanic rocks.* - In Zoubek V. (ed.), Precambrian of Younger Fold Belts. - J. Wiley et Sons, Chichester, 149-152.
- Schreurs J. (1985): Prograde metamorphism of metapelites, garnet-biotite thermometry and prograde changes of biotite chemistry in high grade rocks of West Uusimaa, southwest Finland.* - Lithos, 18, 69-80. Oslo.
- Shaw D.M. (1968): A review of K-Rb fractionation trends by covariance analysis.* - Geochim. cosmochim. Acta, 32, 573-601. Oxford.
- Sighinolfi G.P. (1969): K-Rb ratio in high grade metamor-*

- phism: A confirmation of the hypothesis of a continual crustal evolution. - *Contr. Mineral. Petrol.*, 21, 346-356. Berlin - Heidelberg - New York.
- Sighinolfi G.P. (1971): Investigations into deep crustal levels: Fractionating effects and geochemical trends related to high-grade metamorphism. - Geochim. cosmochim. Acta, 35, 1005-1021. Oxford.*
- Sonder L.J., England P.C., Wemicke B.P. & Christiansen R.L. (1987): A physical model of western North America. In M.P.Coward, J.F.Dewey and P.L.Hancock (eds), Continental Extensional Tectonics, *Geol. Soc. Lond. Spec. Publ.* 28. London.*
- Spear F.S. (1989): Petrologic determination of metamorphic pressure-temperature-time paths. In: F.S.Spear and S.M.Peacock (eds), Metamorphic pressure-temperature-time paths.- Short Course of Geology,7. Amer.Geo-phys.Union, Washington.*
- Spear F.S., Selverstone J., Hickmott D., Crowley P. Hodges K.V. (1984): P-T paths from garnet zoning: a new technique for deciphering tectonic processes in crystalline terranes.- *Geology*, 12, 87-90.*
- Spry A. (1969): Metamorphic textures.- Pergamon, Oxford.*
- Svoboda J. et al. (1966): Regionální geologie ČSSR. Dil I. Český masiv. - Ústř. úst. Geol., Praha.*
- Tamey J. & Windley B.F. (1977): Chemistry, thermal gradients and evolution of the lower continental crust. - *J. Geol. Soc. Lond.*, 134, 153-172. London.*
- Taylor S.R. & McLennan S.M. (1985): The continental crust: Its composition and evolution.- Blackwell Scientific Publ., 312 p. Oxford.*
- Thompson A.B. (1976a): Mineral reactions in pelitic rocks: I. Prediction of P-T-X(Fe-Mg) phase relations.- *Amer.J.Sci.*, 276, 401-424. New Haven.*
- Thompson A.B. (1976): Mineral reactions in pelitic rocks II. Calculations of some P-T-X(Fe-Mg) phase relations.- *Amer. J. Sci.*, 276, 425-444. New Haven.*
- Thompson A.B. (1982): Dehydration melting of pelitic rocks and the generation of H₂O-undersaturated granitic liquid. - *Amer. J. Sci.*, 282, 1567-1595. New Haven.*
- Thompson A.B. (1990): Heat, fluids, and melting in the granulite facies. - In D.Vielzeuf and Ph.Vidal (eds), *Granulites and Crustal Evolution*. - Kluwer Academic Publishers, Netherlands.*
- Tomek Č. & Vrána S. (1992): Deep seismic reflection profiling in the south Bohemian Moldanubium. - Abstract. Styles of Superposed Variscan Nappe Tectonics. 7th Geological workshop - Kutná Hora, ČSAV Praha.*
- Tracy R.J. (1982): Compositional zoning and inclusions in metamorphic minerals. In J.M.Ferry (ed), Characterization of metamorphism through mineral equilibria. *Mineral. Soc. Am. Rev. Mineral.*, 10, 355-397.*
- Tracy R.J., Robinson P. & Thompson A.B. (1976): Garnet composition and zoning in the determination of temperature and pressure of metamorphism, Central Massachusetts. - *Amer. Mineral.*, 61, 762-775.*
- Tuccillo M.E., Essene E.J. & van der Pluijm B.A. (1990): Growth and retrograde zoning in garnets from high-grade metapelites: Implications for pressure-temperature paths. - *Geology*, 18, 839-842. Boulder.*
- Vernon R.H. (1978): Porphyroblast-matrix microstructural relationships in deformed metamorphic rocks. - *Geol. Rdsch.*, 67, 288-305.*
- Vielzeuf D. (1984): Relations des phases dans le facies granulite et implications géodynamiques. L'exemple des granulites des Pyrénées. - *Ann. Sci. Univ. Clermont-Fd.*, 79, Clermont-Ferrand.*
- Vielzeuf D. & Boivin P. (1984): An algorithm for the construction of petrogenetic grids: application to some equilibria in granulitic paragneisses. - *Amer. J.Sci.*, 284, 760-791. New Haven.*
- Vielzeuf D. & Pin C. (1989): Geodynamic implications of granulitic rocks in the Hercynian belt. - *J.S.Daly, R.A.Cliff, and B.W.D. Yardley (eds), Evolution of Metamorphic Belts. - Geol. Soc. Spec. Publ.* 43, 343-348.*
- Vielzeuf D., Clemens J.D., Pin C. & Moinet E. (1990): Granites, granulites and crustal differentiation. - In D.Vielzeuf and Ph.Vidal (eds), *Granulites and Crustal Evolution*. - Kluwer Academic Publishers, Netherlands.*
- Vrána S. (1989): Perpotassic granulites from southern Bohemia - a new rock type derived from partial melting of crustal rocks under upper mantle conditions. - *Contr. Mineral. Petrol.*, 103, 510-522. Berlin - Heidelberg - New York.*
- Vrána S. (1992): The moldanubian zone in southern Bohemia: Polyphase evolution of imbricated crustal and upper mantle segments. In Z.Kukal (ed), *Proceedings of 1st International Conference on the Bohemian Massif.; Geological Survey, Prague (in press)*.*
- Vrána S. & Jakeš P. (1982): Orthopyroxene and two-pyroxene granulites from a segment of charnockite crust in southern Bohemia. - *Věstník Ústř. úst. Geol.*, 57, 129-143, Praha.*
- Webb S.A.C. & Wood B.J. (1986): Spinel-pyroxene-garnet relationships and their dependence on Cr/Al ratio. - *Contr. Mineral. Petrol.*, 92, 471-480. Berlin - Heidelberg - New York.*
- Weber K. & Vollbrecht A. (1989): The crustal structure at the KTB drill site, Oberpfalz. In R.Emmermann and J. Wohlenberg (eds), *The German Continental Deep Drilling Program (KTB). Site-selection studies in the Oberpfalz and Schwarzwald*, 5-36, Springer-Verlag Berlin.*
- Wedepohl K.H. (1991): Chemical composition and fractionation of the continental crust. - *Geol. Rdsch.*, 80, 207-223. Stuttgart.*
- Weisbrod A. (1970): Pétrologie du socle métamorphique des Cévennes médianes (Massif Central français). Reconstitution sédimentologique et approche thermodynamique du métamorphisme. Thèse d'état, Nancy.*
- Wells P.R.A. (1977): Pyroxene thermometry in simple and complex systems. - *Contrib. Mineral. Petrol.*, 62, 129-139. Berlin - Heidelberg - New York.*
- Wendt J.L., Becker H., Fiala J., Kröner A. & Todt W. (in prep.): Evolution of the moldanubian granulite complex, south Bohemia, Czechoslovakia, as revealed by zircon and Sm-Nd dating. In R. Dallmeyer, W. Franke & K.Weber (eds), *Tectonostratigraphic evolution of central European Paleozoic orogenes*. Springer-Verlag, Berlin.*

- Winkler H.G.F. & von Platen H. (1961): Experimentelle Gesteinsmetamorphose. - Geochim. cosmochim. Acta, 24, 48-69. Oxford.*
- Wood B.J. & Banno S. (1973): Garnet-orthopyroxene and orthopyroxene-clinopyroxene relationships in simple and complex systems. - Contr. Mineral. Petrol., 42, 109-124. Berlin - Heidelberg - New York.*
- Yardley B.W.D. (1977): An empirical study of diffusion in garnets. - Amer. Mineral., 62, 793-800. Washington.*
- Zartner W.R. (1929): Geologisch-petrographische studien im Egertalgebiet zwischen Warta und Kaaden. - Sbor. Stát. geol. Úst. ČSR, VIII, 105-188. Praha.*
- Zoubek V. (1948a): Poznámky k otázce skarnů, granulitů a jihočeských grafitových ložisek. - Sbor. Stát. geol. Úst., 13, 483-498. Praha.*
- Zoubek V. (1948b): Poznámky ke geologii krystalinika Českého masivu. - Sbor. Stát. geol. Úst., 15, 339-398. Praha.*
- Zoubek V. & Škvor V. (eds) (1963): Vysvětlivky k přehledné geologické mapě 1:200 000 M-33-XIV Teplice, M-33-VIII Chabařovice. - Ústř. úst. Geol., Nakl. ČSAV, Praha.*
- Zoubek V. et al. (1988): Precambrian of Younger Fold Belts. - J. Wiley et Sons, Chichester, 885 pp.*